Neuroscience Letters, 154 (1993) 218–220
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, K., see Aoki, M., 69
Acosta, G.B., Losada, M.E.O. and Rubio, M.C., Area-dependent changes in GABAergic function after acute and chronic cold stress, 175
Akiguchi, I., see Nakamura, S., 61
Alonzo, G., see Andrews, R.J., 9
Anderson, T., Schwarcz, R., Löve, A. and Kristensson, K., Measles virus-induced hippocampal neurodegeneration in the mouse: a novel, subacute model for testing neuroprotective agents, 109
Andrews, R.J., Bringas, J.R., Alonzo, G., Salamat, M.S., Khoshyomn, S. and Gluck, D.S., Corpus callosotomy effects on cerebral blood flow and evoked potentials (transcallosal diaschisis), 9
Aoki, M., Abe, K., Liu, X.-H., Lee, T.-H., Kato, H. and Kogure, K., Reduction of HSP70 and HSC70 mRNA inductions by bifemelane hydrochloride after transient ischemia in gerbil brain, 69
Austin, D.R., see Gold, B.G., 129
Avoli, M., see Mattia, D., 157

Bandler, R., see Keay, K.A., 23
Barnes, D., see Weisz, P.V., 153
Behl, C. and Schubert, D., Heat shock partially protects rat pheochromocytoma PC12 cells from amyloid β peptide toxicity, 1
Benedikz, E., see Casaccia-Bonnefil, P., 5
Berezin, V.A., see Krivko, I.M., 17
Berger, R.J., Cooling down to hibernate: sleep and hibernation constitute a physiological continuum of energy conservation, 213
Bergeron, C., see Morrison, L.D., 141
Bergold, P.J., see Casaccia-Bonnefil, P., 5
Bhargava, S. and Rao, P.D.P., Distribution of corticotropin-releasing factor immunoreactive neurons in the brain of the tigerfrog, *Rana tigrina*, 27
Bijak, M., see Jarolimek, W., 31
Billard, J.M., see Deisz, R.A., 209
Bohus, B., see Krugers, H.J., 73
Bollensen, E., see Zahn, M., 93
Braak, E. and Braak, H., The new monodendritic neuronal type within the adult human cerebellar granule cell layer shows calretinin-immunoreactivity, 199
Braak, H., see Braak, E., 199
Bringas, J.R., see Andrews, R.J., 9
Brown, L.M., Developmental alterations in *N*-methyl-D-aspartate stimulated [^{3}H]norepinephrine release in rat brain cortex and hippocampus, 43
Burwell, R.D., see Whealin, J.M., 149

Candy, J.M., see Gocht, A., 187
Casaccia-Bonnefil, P., Benedikz, E., Rai, R. and Bergold, P.J., Excitatory and inhibitory pathways modulate kainate excitotoxicity in hippocampal slice cultures, 5
Chartier-Harlin, M.-C., see Parfitt, M., 81
Confort-Gouny, S., see Nicoli, F., 47
Cozzone, P.J., see Nicoli, F., 47
Crawford, F., see Houlden, H., 161
Crook, R., see Parfitt, M., 81

Dang, N., see Fukuchi, K.-i., 145
Davies, H.A., see Rusakov, D.A., 13
Deisz, R.A., Billard, J.M. and Zieglgänsberger, W., Pre- and postsynaptic $GABA_B$ receptors of rat neocortical neurons differ in their pharmacological properties, 209
Delwaide, C., see Nicoli, F., 47
Demmelhuber, J., see Jarolimek, W., 31
Dermietzel, R., see Rohlmann, A., 206
Dieringer, N., see Reichenberger, I., 89
Dirnagl, U., see Villringer, A., 101
Dobie, D.J., see Merchant, K.M., 163
Dobo, E., see Rohlmann, A., 206
Dorsa, D.M., see Merchant, K.M., 163
Drucker-Colín, R., see Prospero-García, O., 168
Dykstra, C., see Meller, S.T., 137

Efthimiopoulos, S., see Shioi, J., 121
Endoh, M., Maiese, K., Pulsinellid, W.A. and Wagner, J.A., Reactive astrocytes express NADPH diaphorase in vivo after transient ischemia, 125

Felgenhauer, K., see Zahn, M., 93
Fukuchi, K.-i., Sopher, B., Furlong, C.E., Smith, A.C., Dang, N. and Martin, G.M., Selective neurotoxicity of COOH-terminal fragments of the β-amyloid precursor protein, 145
Fukunaga, R., see Yamaoka, Y., 20
Furlong, C.E., see Fukuchi, K.-i., 145

Gallagher, M., see Whealin, J.M., 149
Gebhart, G.F., see Meller, S.T., 137
Geisert Jr., E.E., see Irwin, M.H., 57
Gluck, D.S., see Andrews, R.J., 9
Gocht, A., Keith, A.B., Candy, J.M. and Morris, C.M., Iron uptake in the brain of the myelin-deficient rat, 187
Gold, B.G., Storm-Dickerson, T. and Austin, D.R., Regulation of the transcription factor c-JUN by nerve growth factor in adult sensory neurons, 129
Goldsmith, S., see Joyce, J.N., 171
Grover, L.M. and Teyler, T.J., Role of adenosine in heterosynaptic, posttetanic depression in area CA1 of hippocampus, 39

Harrison, E., see Rusakov, D.A., 13
Hock, C., see Villringer, A., 101
Hofer, A., see Rohlmann, A., 206
Houlden, H., Crawford, F., Rossor, M. and Mullan, M., Screening for the APP codon 670/671 mutations in Alzheimer's disease, 161
Hwa, G.G.C., see Mattia, D., 157

Irwin, M.H. and Geisert Jr., E.E., The upregulation of a glial cell surface antigen at the astrocytic scar in the rat, 57
Ito, H., see Uchiyama, H., 35
Iwamoto, Y., see Kamei, J., 65

Jarolimek, W., Demmelhuber, J., Bijak, M. and Misgeld, U., CGP 55845A blocks baclofen, γ-aminobutyric acid and inhibitory postsynaptic potassium currents in guinea pig CA3 neurons, 31
Jiménez-Anguiano, A., see Prospero-García, O., 168
Joyce, J.N., Kaeger, C., Ryoo, H. and Goldsmith, S., Dopamine D2 receptors in the hippocampus and amygdala in Alzheimer's disease, 171

Juusola, M. and Weckström, M., Band-pass filtering by voltage-dependent membrane in an insect photoreceptor, 84

Kaeger, C., see Joyce, J.N., 171
Kamei, J., Iwamoto, Y., Suzuki, T., Misawa, M., Nagase, H. and Kasuya, Y., Antitussive effects of μ- and κ-agonists in diabetic rats, 65
Kasuya, Y., see Kamei, J., 65
Kato, H., see Aoki, M., 69
Keay, K.A. and Bandler, R., Deep and superficial noxious stimulation increases Fos-like immunoreactivity in different regions of the midbrain periaqueductal grey of the rat, 23
Keith, A.B., see Gocht, A., 187
Khoshyomn, S., see Andrews, R.J., 9
Kimura, J., see Nakamura, S., 61
Kimura, J., see Yamaoka, Y., 20
Kish, S.J., see Morrison, L.D., 141
Kogure, K., see Aoki, M., 69
Kononenko, N.I., see Storozhuk, M.V., 203
Koolhaas, J.M., see Krugers, H.J., 73
Korf, J., see Krugers, H.J., 73
Kostyuk, P.G., see Storozhuk, M.V., 203
Kothbauer, K., Schmid, U.D., Liechti, S. and Rösler, K.M., The effect of ketamine anesthetic induction on muscle responses to transcranial magnetic cortex stimulation studied in man, 105
Kristensson, K., see Anderson, T., 109
Krivko, I.M., Rusakov, D.A., Savina, S.V., Skibo, G.G. and Berezin, V.A., Changes in the neural cell adhesion molecule patterns on the rat glial cell surfaces with development and contact formation in vitro, 17
Krugers, H.J., Koolhaas, J.M., Bohus, B. and Korf, J., A single social stress-experience alters glutamate receptor-binding in rat hippocampal CA3 area, 73

Laskawi, R., see Rohlmann, A., 206
Lee, T.-H., see Aoki, M., 69
Leung, L.S., see Zhao, D., 191
Li Volsi, G., see Licata, F., 195
Licata, F., Li Volsi, G., Maugeri, G. and Santangelo, F., Excitatory and inhibitory effects of 5-hydroxytryptamine on the firing rate of medial vestibular nucleus neurons in the rat, 195
Liechti, S., see Kothbauer, K., 105
Liu, X.-H., see Aoki, M., 69
Losada, M.E.O., see Acosta, G.B., 175
Löve, A., see Anderson, T., 109

Mäder, M., see Zahn, M., 93
Maiese, K., see Endoh, M., 125
Maloteaux, J.M., see Nicoli, F., 47
Martin, G.M., see Fukuchi, K.-i., 145
Maruyama, S., see Sasaki, S., 117
Maruyama, W., see Ota, A., 183
Mason, P., Lipopolysaccharide induces fever and decreases tail flick latency in awake rats, 134
Mastaglia, F.L., see Wilson, S.A., 52
Mattia, D., Hwa, G.G.C. and Avoli, M., Epileptiform activity induced by 4-aminopyridine in guinea-pig and rat neocortices, 157
Maugeri, G., see Licata, F., 195
Maves, T.J., see Meller, S.T., 137
Meller, S.T., Dykstra, C., Pechman, P.S., Maves, T.J. and Gebhart, G.F., Ethanol dose-dependently attenuates NMDA-mediated thermal hyperalgesia in the rat, 137
Merchant, K.M., Dobie, D.J. and Dorsa, D.M., Differential loss of dopamine D2 receptor mRNA isoforms during aging in Fischer-344 rats, 163
Misawa, M., see Kamei, J., 65
Misgeld, U., see Jarolimek, W., 31
Morris, C.M., see Gocht, A., 187
Morrison, L.D., Bergeron, C. and Kish, S.J., Brain *S*-adenosylmethionine decarboxylase activity is increased in Alzheimer's disease, 141
Mullan, M., see Houlden, H., 161

Nagase, H., see Kamei, J., 65
Nagatsu, T., see Ota, A., 183
Nakamura, S., Akiguchi, I. and Kimura, J., The histochemical demonstration of MPTP oxidation in the postmortem human striatum, 61
Naoi, M., see Ota, A., 183
Nicoli, F., Vion-Dury, J., Maloteaux, J.M., Delwaide, C., Confort-Gouny, S., Sciaky, M. and Cozzone, P.J., CSF and serum metabolic profile of patients with Huntington's chorea: a study by high resolution proton NMR spectroscopy and HPLC, 47

Ohno-Shosaku, T., see Yamamoto, C., 179
Ota, A., Maruyama, W., Takahashi, T., Naoi, M. and Nagatsu, T., The effect of a heterocyclic amine, 3-amino-1,4-dimethyl-5*H*-pyrido [4,3-*b*]indole (Trp-P-1), on tyrosine hydroxylase mRNA in PC12h cells, 183
Ottersen, O.P., see Reichenberger, I., 89

Paré, D. and Steriade, M., The reticular thalamic nucleus projects to the contralateral dorsal thalamus in macaque monkey, 96
Parfitt, M., Crook, R., Roques, P., Rossor, M. and Chartier-Harlin, M.-C., The Cystatin-C gene is not linked to early onset familial Alzheimer's disease, 81
Pechman, P.S., see Meller, S.T., 137
Planck, J., see Villringer, A., 101
Porter, R.H.P. and Roberts, P.J., Glutamate metabotropic receptor activation in neonatal rat cerebral cortex by sulphur-containing excitatory amino acids, 78
Prospero-García, O., Jiménez-Anguiano, A. and Drucker-Colín, R., Chloramphenicol prevents carbachol-induced REM sleep in cats, 168
Pulsinellid, W.A., see Endoh, M., 125

Rai, R., see Casaccia-Bonnefil, P., 5
Rao, P.D.P., see Bhargava, S., 27
Reeh, P.W., see Steen, K.H., 113
Refolo, L.M., see Shioi, J., 121
Reichenberger, I., Streit, P., Ottersen, O.P. and Dieringer, N., GABA- and glycine-like immunoreactivities in the cerebellum of the frog, 89
Rösler, K.M., see Kothbauer, K., 105
Robakis, N.K., see Shioi, J., 121
Roberts, P.J., see Porter, R.H.P., 78
Rohlmann, A., Laskawi, R., Hofer, A., Dobo, E., Dermietzel, R. and Wolff, J.R., Facial nerve lesions lead to increased immunostaining of the astrocytic gap junction protein (connexin 43) in the corresponding facial nucleus of rats, 206
Roques, P., see Parfitt, M., 81
Rossor, M., see Houlden, H., 161
Rossor, M., see Parfitt, M., 81
Rubio, M.C., see Acosta, G.B., 175
Rusakov, D.A., Stewart, M.G., Davies, H.A. and Harrison, E., Spatial

re-arrangement of the vesicle apparatus in forebrain synapses of chicks 30 min after passive avoidance training, 13
Rusakov, D.A., see Krivko, I.M., 17
Ryoo, H., see Joyce, J.N., 171

Salamat, M.S., see Andrews, R.J., 9
Santangelo, F., see Licata, F., 195
Sasaki, S. and Maruyama, S., Ultrastructural study of Bunina bodies in the anterior horn neurons of patients with amyotrophic lateral sclerosis, 117
Savina, S.V., see Krivko, I.M., 17
Sawada, S., see Yamamoto, C., 179
Schleinkofer, L., see Villringer, A., 101
Schmid, U.D., see Kothbauer, K., 105
Schmidt, B., see Zahn, M., 93
Schubert, D., see Behl, C., 1
Schwarcz, R., see Anderson, T., 109
Sciaky, M., see Nicoli, F., 47
Shimohama, S., see Yamaoka, Y., 20
Shioi, J., Refolo, L.M., Efthimiopoulos, S. and Robakis, N.K., Chondroitin sulfate proteoglycan form of cellular and cell-surface Alzheimer amyloid precursor, 121
Skibo, G.G., see Krivko, I.M., 17
Smith, A.C., see Fukuchi, K.-i., 145
Solem, M., see Weisz, P.V., 153
Sopher, B., see Fukuchi, K.-i., 145
Steen, K.H. and Reeh, P.W., Sustained graded pain and hyperalgesia from harmless experimental tissue acidosis in human skin, 113
Steriade, M., see Paré, D., 96
Stewart, M.G., see Rusakov, D.A., 13
Storm-Dickerson, T., see Gold, B.G., 129
Storozhuk, M.V., Kostyuk, P.G. and Kononenko, N.I., Patch-clamp recording of cAMP-induced membrane current noise in *Helix pomatia* neurons, 203
Streit, P., see Reichenberger, I., 89
Suzuki, T., see Kamei, J., 65

Takahashi, T., see Ota, A., 183
Taniguchi, T., see Yamaoka, Y., 20
Teyler, T.J., see Grover, L.M., 39
Thickbroom, G.W., see Wilson, S.A., 52

Uchiyama, H. and Ito, H., Target cells for the isthmo-optic fibers in the retina of the Japanese quail, 35

Villringer, A., Planck, J., Hock, C., Schleinkofer, L. and Dirnagl, U., Near infrared spectroscopy (NIRS): a new tool to study hemodynamic changes during activation of brain function in human adults, 101
Vion-Dury, J., see Nicoli, F., 47

Wagner, J.A., see Endoh, M., 125
Weckström, M., see Juusola, M., 84
Weisz, P.V., Solem, M. and Barnes, D., Expression of a TGFβ regulated, brain-specific mRNA in serum-free mouse embryo (SFME) cells, 153
Whealin, J.M., Burwell, R.D. and Gallagher, M., The effects of aging on diurnal water intake and melatonin binding in the suprachiasmatic nucleus, 149
Wilson, S.A., Thickbroom, G.W. and Mastaglia, F.L., Topography of excitatory and inhibitory muscle responses evoked by transcranial magnetic stimulation in the human motor cortex, 52
Wolff, J.R., see Rohlmann, A., 206

Yamamoto, C., Sawada, S. and Ohno-Shosaku, T., Electrophysiological features of habenular neurons of the guinea pig: high efficacy of synaptic transmission and variable configuration of unitary spikes, 179
Yamaoka, Y., Shimohama, S., Kimura, J., Fukunaga, R. and Taniguchi, T., Changes in protein kinase C isozymes in the rat hippocampus following transient hypoxia, 20

Zahn, M., Mäder, M., Schmidt, B., Bollensen, E. and Felgenhauer, K., Purification and N-terminal sequence of β-trace, a protein abundant in human cerebrospinal fluid, 93
Zhao, D. and Leung, L.S., Partial hippocampal kindling increases paired-pulse facilitation and burst frequency in hippocampal CA1 neurons, 191
Zieglgänsberger, W., see Deisz, R.A., 209